arXiv:1402.2856v2 [math.DG] 3 Apr 2014

# A FAMILY OF MAPS WITH MANY SMALL FIBERS

HANNAH ALPERT AND LARRY GUTH

Abstract. The waist inequality states that for a continuous map from $S^n$ to $\mathbb{R}^q$, not all fibers can have small $(n-q)$-dimensional volume. We construct maps for which most fibers have small $(n-q)$-dimensional volume and all fibers have bounded $(n-q)$-dimensional volume.

Let $n, q \in \mathbb{N}$ with $n > q \geq 1$, and let $f : S^n \to \mathbb{R}^q$ be a continuous map. Let $\widehat{p} : \mathbb{R}^{n+1} \to \mathbb{R}^q$ be a surjective linear map, and let $p = \widehat{p}|_{S^n}$. The waist inequality states that the largest fiber of $f$ is at least as large as the largest fiber of $p$:

$$\sup_{y \in \mathbb{R}^q} \mathrm{Vol}_{n-q} f^{-1}(y) \geq \sup_{y \in \mathbb{R}^q} \mathrm{Vol}_{n-q} p^{-1}(y).$$

See [1], [3], [4], and [6] for proofs of the waist inequality, or [5] for a survey. In the case $q = 1$, the waist inequality is a consequence of the isoperimetric inequality on $S^n$. The isoperimetric inequality can also be used to prove that the portion of $S^n$ covered by small fibers of $f$ is not very big; that is, for all $\varepsilon$, we have

$$\mathrm{Vol}_n \ f^{-1}\{y : \mathrm{Vol}_{n-q} f^{-1}(y) < \varepsilon\} \leq \mathrm{Vol}_n \ p^{-1}\{y : \mathrm{Vol}_{n-q} p^{-1}(y) < \varepsilon\}.$$

The theorem presented in this paper describes how the same statement does not hold in the case $q > 1$. We have also included an appendix with a more precise statement of the waist inequality and the isoperimetric inequality.

**Theorem 1.** *For every $n, q \in \mathbb{N}$ with $n > q > 1$, and for every $\varepsilon > 0$, there is a continuous map $f : S^n \to \mathbb{R}^q$ such that all but $\varepsilon$ of the $n$-dimensional volume of $S^n$ is covered by fibers that have $(n-q)$-dimensional volume at most $\varepsilon$. Moreover, we may require that every fiber of $f$ has $(n-q)$-dimensional volume bounded by $C_{n,q}$, a constant not depending on $\varepsilon$.*

In what follows, $I^n = [0,1]^n$ denotes the $n$-dimensional unit cube, and $\partial I^n$ denotes its boundary. A *tree* refers to the topological space corresponding to a graph-theoretic tree: topologically, a tree is a finite 1-dimensional simplicial complex that is contractible.

The bulk of the construction comes from the following lemma, in which we construct a preliminary "tree map" $t_{n,r,\delta}$ from $I^n$ to a tree. Later, to construct $f$ we will change the domain from $I^n$ to $S^n$ by gluing several tree maps together, and we will change the range from the tree to $\mathbb{R}^q$ by composing with a map from a thickened tree to $\mathbb{R}^q$. In the tree map $t_{n,r,\delta}$, the parameter $r$ corresponds to the depth of the tree. As $r$ increases, the typical fiber of the map becomes smaller. The parameter $\delta$ corresponds to the total volume of the larger fibers.

**Lemma 1.** *For every $n, r \in \mathbb{N}$, there is a rooted tree $T_{n,r}$ such that for every $\delta > 0$ there is a continuous map $t_{n,r,\delta} : I^n \to T_{n,r}$ with the following properties:*

2010 *Mathematics Subject Classification.* 53C23.



FIGURE 1. Every fiber of $t_{2,2,\delta}$ has length at most 6, and most fibers have length at most 1.

(1) *Every fiber of* $t_{n,r,\delta}$ *is either a single point, the boundary of an* $n$*-dimensional cube of side length at most* 1*, or the* $(n-1)$*-skeleton of a* $2 \times 2 \times \cdots \times 2$ *array of* $n$*-dimensional cubes each of side length at most* $\frac{1}{2}$*.*
(2) *All but* $\delta$ *of the volume of* $I^n$ *is covered by fibers of* $t_{n,r,\delta}$ *that are boundaries of* $n$*-dimensional cubes of side length at most* $2^{-r}$*.*
(3) $t_{n,r,\delta}(\partial I^n)$ *is a single point, the root of* $T_{n,r}$*.*
(4) *Each vertex has at most* $2^n$ *daughter vertices.*

*Proof.* We construct the tree and tree map recursively in $r$. For $r = 0$, the tree $T_{n,0}$ is a single edge which we may identify with the interval $[0, \frac{1}{2}]$, with 0 being the root. For any $\delta$, we set $t_{n,0,\delta}(x) = \text{dist}(x, \partial I^n)$ for all $x \in I^n$.

Now let $r > 0$. To construct $T_{n,r}$, we take the disjoint union of one copy of $[0, 1]$ and $2^n$ copies of $T_{n,r-1}$, and identify the root of every copy of $T_{n,r-1}$ with $1 \in [0, 1]$. The root of $T_{n,r}$ is $0 \in [0, 1]$. We define $t_{n,r,\delta}$ piecewise as follows. For some small choice of $\delta_1 > 0$, we define $t_{n,r,\delta}$ on the closed $\delta_1$-neighborhood of $\partial I^n$ to $[0, 1] \subset T_{n,r}$ by

$$t_{n,r,\delta}(x) = \frac{1}{\delta_1}\text{dist}(x, \partial I^n).$$

Then, translating the coordinate hyperplanes to pass through the center of $I^n$ we divide the remainder of the cube into a $2 \times 2 \times \cdots \times 2$ array of cubes $Q_1, \ldots, Q_{2^n}$ each of side length slightly less than $\frac{1}{2}$. For each $j = 1, \ldots 2^n$, let $\lambda_j : Q_j \to I^n$ be the map that scales $Q_j$ up to unit size, and let $i_j : T_{n,r-1} \to T_{n,r}$ be the inclusion of the $j$th copy of $T_{n,r-1}$ into $T_{n,r}$. Then for some small choice of $\delta_2 > 0$, we put

$$t_{n,r,\delta}|_{Q_j} = i_j \circ t_{n,r-1,\delta_2} \circ \lambda_j.$$

Properties 1, 3, and 4 are easily satisfied by the construction. To ensure property 2, we need to choose $\delta_1$ and $\delta_2$. The volume of $I^n$ that is covered by large fibers—fibers not equal to the boundary of a cube of side length at most $2^{-r}$—is at most $\delta_1 \cdot 2n + 2^n \cdot \delta_2 \cdot 2^{-n}$, because the area of $\partial I^n$ is $2n$ and because the portion of each $Q_j$ that is covered by large fibers has volume at most $\delta_2 \cdot \mathrm{Vol}(Q_j) < \delta_2 \cdot 2^{-n}$. Thus we may choose $\delta_1 = \frac{\delta}{4n}$ and $\delta_2 = \frac{\delta}{2}$. □

*Proof of Theorem 1.* We may replace $S^n$ by $\partial I^{n+1}$ by composing with the (bi-Lipschitz) homeomorphism $\psi : S^n \to \partial I^{n+1}$ given by lining up the centers of $S^n$ and $\partial I^{n+1}$ in $\mathbb{R}^{n+1}$ and projecting radially. We start by constructing a tree $T$ and a tree map $t : \partial I^{n+1} \to T$. For some large choice of $r$, let $T$ be the tree obtained by identifying the roots of $2(n+1)$ copies of $T_{n,r}$, one for each $n$-dimensional face of $\partial I^{n+1}$. For some small choice of $\delta$, define $t$ on each $n$-dimensional face of $\partial I^{n+1}$ to be the composition of $t_{n,r,\delta}$ with the inclusion of the corresponding $T_{n,r}$ into $T$.

The fibers of $t$ have dimension $n-1$. In order to cut the fibers down to dimension $n-q$, we next construct a projection map $p : \partial I^{n+1} \to \mathbb{R}^{q-1}$ such that the fibers of $p$ intersect the fibers of $t$ transversely. The fibers of $t$ have codimension 2 in $\mathbb{R}^{n+1}$ and are aligned with the standard coordinates, so we achieve transversality by using other linear coordinates to construct $p$. We choose $q-1$ linearly independent vectors $v_1, \ldots, v_{q-1} \in \mathbb{R}^{n+1}$ such that for every two standard basis vectors $e_i, e_j \in \mathbb{R}^{n+1}$ the spaces $\mathrm{span}\{e_i, e_j\}^\perp$ and $\mathrm{span}\{v_1, \ldots, v_{q-1}\}^\perp$ intersect transversely; equivalently, the set $e_i, e_j, v_1, \ldots, v_{q-1}$ is linearly independent. For $k = 1, \ldots, q-1$, define the $k$th component of $p$ to be the dot product of the input with $v_k$. Then the fibers of $t \times p : \partial I^{n+1} \to T \times \mathbb{R}^{q-1}$ are codimension $q-1$ transverse linear cross-sections of the $(n-1)$-dimensional fibers of $t$, and have $(n-q)$-dimensional volume bounded by some constant depending on $n$ and $q$.

There exists $M$ large enough that $p(\partial I^{n+1})$ is contained in the $(q-1)$-dimensional ball $B(M)$ of radius $M$. We define a map $\phi : T \times B(M) \to \mathbb{R}^q$ such that the number of points in each fiber of $\phi$ is at most the maximum degree of $T$, which is $2^n + 1$. Then we define $f = \phi \circ (t \times p)$. The fibers of $f$, like the fibers of $t \times p$, have $(n-q)$-dimensional volume bounded by a constant $C_{n,q}$.

The map $\phi$ is constructed as follows. Let $\phi|_{T \times \{0\}}$ be an embedding of $T$ into $\mathbb{R}^q$ in which the edges map to straight line segments and each daughter vertex has $x_1$-coordinate greater than that of its parent. Let $d$ be the minimum distance between disjoint edges of $\phi(T \times \{0\})$. Then for every $p \in T$ and $x \in B(M)$, we set

$$\phi(p, x) = \phi(p, 0) + \frac{d}{4}\left(0, \frac{x}{M}\right),$$

where $\left(0, \frac{x}{M}\right)$ denotes the point in $\mathbb{R}^q$ constructed by adding onto $\frac{x}{M} \in \mathbb{R}^{q-1}$ a first coordinate of 0. If $\phi(p, x) = \phi(p', x')$, then $\phi(p, 0)$ and $\phi(p', 0)$ are at most $\frac{d}{2}$ apart, so $p$ and $p'$ lie on two incident edges of $T$; also, $\phi(p, 0)$ and $\phi(p', 0)$ have the same

$x_1$-coordinate, so these two edges are between two daughters and a common parent, rather than a daughter, a parent, and a grandparent.

To finish the proof, we show that $\delta$ and $r$ may be chosen such that all but $\varepsilon$ of the $n$-dimensional volume of $\partial I^{n+1}$ is covered by fibers with $(n-q)$-dimensional volume at most $\varepsilon$. The maximum number of daughter vertices of any vertex of $T$ is $2^n$, and most of $\partial I^{n+1}$ is covered by fibers of $f$ that are unions of at most $2^n$ codimension $q-1$ transverse linear cross-sections of boundaries of $n$-dimensional cubes of side length at most $2^{-r}$. We choose $r$ large enough that every codimension $q-1$ transverse linear cross-section of $2^{-r}\partial I^n$ has $(n-q)$-dimensional volume at most $\frac{\varepsilon}{2^n}$. The volume of the portion of $\partial I^{n+1}$ covered by larger fibers is at most $2(n+1)\cdot\delta$, so we choose $\delta < \frac{\varepsilon}{2(n+1)}$. □

## Appendix: The waist inequality and the isoperimetric inequality

In order to be precise about the waist inequality, we need a notion of $(n-q)$-dimensional volume of arbitrary closed subsets in $S^n$. Gromov's version of the waist inequality is stated in terms of the Lebesgue measures $\mathrm{Vol}_n$ of the $\varepsilon$-neighborhoods $f^{-1}(y)_\varepsilon$ of the fibers $f^{-1}(y)$ of a continuous map $f$.

**Theorem 2** (Waist inequality, [4])**.** *Let $f : S^n \to \mathbb{R}^q$ be a continuous map. Then there exists a point $y \in \mathbb{R}^q$ such that for all $\varepsilon > 0$, we have*

$$\mathrm{Vol}_n(f^{-1}(y)_\varepsilon) \geq \mathrm{Vol}_n(S^{n-q}_\varepsilon),$$

*where $S^{n-q} \subset S^n$ denotes an equatorial $(n-q)$-sphere.*

The paper [6] gives a detailed exposition of the proof of the waist inequality and fills in some small gaps in the original argument. For convenience we introduce a notation for comparing the $\varepsilon$-neighborhoods of two sets: given $E, F \subseteq S^n$, we say that $E$ is ***larger in neighborhood*** than $F$, denoted $E \geq_{\mathrm{nbd}} F$, if for all $\varepsilon > 0$ we have

$$\mathrm{Vol}_n(E_\varepsilon) \geq \mathrm{Vol}_n(F_\varepsilon).$$

Then the waist inequality states that for some $y \in \mathbb{R}^q$ we have $f^{-1}(y) \geq_{\mathrm{nbd}} S^{n-q}$.

In the case $q = 1$, we would like to say that the waist inequality is a consequence of the isoperimetric inequality. The classical isoperimetric inequality applies only to regions with smooth boundary, so we need the following version, which is stated and proved in [2] and attributed to [7]:

**Theorem 3** (Isoperimetric inequality)**.** *Let $A \subseteq S^n$ be a closed set and $B \subseteq S^n$ be a closed ball with $\mathrm{Vol}_n(B) = \mathrm{Vol}_n(A)$. Then we have*

$$A \geq_{\mathrm{nbd}} B.$$

In the introduction we claimed that in the case $q = 1$, the isoperimetric inequality could be used to prove, in addition to the waist inequality, another statement about the volume of $S^n$ covered by small fibers. Here we formulate the statement more precisely and prove it. The proof implies the waist inequality for $q = 1$.

**Theorem 4.** *Let $f : S^n \to \mathbb{R}$ be a continuous map, and $p : S^n \to \mathbb{R}$ be the restriction to $S^n$ of a surjective linear map $\widehat{p} : \mathbb{R}^{n+1} \to \mathbb{R}$. Then for all $y \in p(S^n)$, we have*

$$\mathrm{Vol}_n\{x \in S^n : f^{-1}(f(x)) \geq_{\mathrm{nbd}} p^{-1}(y)\} \geq \mathrm{Vol}_n\{x \in S^n : p^{-1}(p(x)) \geq_{\mathrm{nbd}} p^{-1}(y)\}.$$

The proof of this theorem is based on the following lemma:

**Lemma 2.** *Let $X, Y \subset S^n$ be closed sets with $X \cup Y = S^n$. Let $B^X, B^Y \subset S^n$ be closed balls such that their two centers are antipodal in $S^n$ and $\mathrm{Vol}_n(B^X) = \mathrm{Vol}_n(X)$ and $\mathrm{Vol}_n(B^Y) = \mathrm{Vol}_n(Y)$. Then we have*

$$X \cap Y \geq_{\mathrm{nbd}} B^X \cap B^Y.$$

*Proof.* First we claim that $(X \cap Y)_\varepsilon$ is the disjoint union of $X_\varepsilon \setminus X$, $Y_\varepsilon \setminus Y$, and $X \cap Y$. It is clear that $(X \cap Y)_\varepsilon$ is the disjoint union of its intersections with $S^n \setminus X$, $S^n \setminus Y$, and $X \cap Y$. Thus it suffices to show that

$$(X \cap Y)_\varepsilon \cap (S^n \setminus X) = X_\varepsilon \setminus X.$$

Because $(X \cap Y)_\varepsilon \subseteq X_\varepsilon$, we immediately have

$$(X \cap Y)_\varepsilon \cap (S^n \setminus X) \subseteq X_\varepsilon \setminus X.$$

For the reverse inclusion, let $y \in X_\varepsilon \setminus X$, and let $\gamma : [0,1] \to S^n$ be a curve of length at most $\varepsilon$ with $\gamma(0) = y$ and $\gamma(1) = x \in X$. Let $t \in [0,1]$ be the greatest value with $\gamma(t) \in Y$. Then $\gamma(t) \in X \cap Y$, so $y \in (X \cap Y)_\varepsilon$.

Thus, applying the isoperimetric inequality and additivity of measure, we have

$$\begin{aligned} \mathrm{Vol}_n((X \cap Y)_\varepsilon) &= \mathrm{Vol}_n(X_\varepsilon) - \mathrm{Vol}_n(X) + \mathrm{Vol}_n(Y_\varepsilon) - \mathrm{Vol}_n(Y) + \mathrm{Vol}_n(X \cap Y) \geq \\ &\geq \mathrm{Vol}_n(B^X_\varepsilon) - \mathrm{Vol}_n(B^X) + \mathrm{Vol}_n(B^Y_\varepsilon) - \mathrm{Vol}_n(B^Y) + \mathrm{Vol}_n(B^X \cap B^Y) = \\ &= \mathrm{Vol}_n((B^X \cap B^Y)_\varepsilon). \end{aligned}$$

□

*Proof of Theorem 4.* Without loss of generality we assume $p(S^n) = [0,1]$ and $y \leq \frac{1}{2}$. Then on the right-hand side of the desired inequality we have

$$\{x \in S^n : p^{-1}(p(x)) \geq_{\mathrm{nbd}} p^{-1}(y)\} = p^{-1}[y, 1-y].$$

Define $\alpha, \beta \in \mathbb{R}$ as

$$\alpha = \sup\{t \in \mathbb{R} : \mathrm{Vol}_n f^{-1}(-\infty, t) \leq \mathrm{Vol}_n p^{-1}[0, y]\},$$

$$\beta = \inf\{t \in \mathbb{R} : \mathrm{Vol}_n f^{-1}(t, \infty) \leq \mathrm{Vol}_n p^{-1}[y, 1]\}.$$

For each $t \in [\alpha, \beta]$, apply the lemma with $X = f^{-1}(-\infty, t]$ and $Y = f^{-1}[t, \infty)$ to get $f^{-1}(t) \geq_{\mathrm{nbd}} p^{-1}[y_1, y_2]$ for some $y_1, y_2 \in [y, 1-y]$. In particular, we have

$$f^{-1}(t) \geq_{\mathrm{nbd}} p^{-1}(y_1) \geq_{\mathrm{nbd}} p^{-1}(y).$$

Thus, we have

$$f^{-1}[\alpha, \beta] \subseteq \{x \in S^n : f^{-1}(f(x)) \geq_{\mathrm{nbd}} p^{-1}(y)\}.$$

Because $\mathrm{Vol}_n f^{-1}(-\infty, \alpha) \leq \mathrm{Vol}_n p^{-1}[0, y]$ and $\mathrm{Vol}_n f^{-1}(\beta, \infty) \leq \mathrm{Vol}_n p^{-1}[y, 1]$ we have

$$\mathrm{Vol}_n f^{-1}[\alpha, \beta] \geq \mathrm{Vol}_n p^{-1}[y, 1-y].$$

□

## References

[1] F.J. Almgren, *The theory of varifolds — a variational calculus in the large for the k-dimensional area integrated*, Mimeographed notes, 1965.

[2] T. Figiel, J. Lindenstrauss, and V. D. Milman, *The dimension of almost spherical sections of convex bodies*, Acta Math. **139** (1977), no. 1-2, 53–94. MR 0445274 (56 #3618)

[3] M. Gromov, *Filling Riemannian manifolds*, J. Differential Geom. **18** (1983), no. 1, 1–147. MR 697984 (85h:53029)

[4] ———, *Isoperimetry of waists and concentration of maps*, Geom. Funct. Anal. **13** (2003), no. 1, 178–215. MR 1978494 (2004m:53073)

[5] L. Guth, *The waist inequality in Gromov's work*, The Abel Prize 2008–2012 (H. Holden and R. Piene, eds.), Springer, 2014, pp. 181–195.

[6] Y. Memarian, *On Gromov's waist of the sphere theorem*, J. Topol. Anal. **3** (2011), no. 1, 7–36. MR 2784762 (2012g:53066)

[7] E. Schmidt, *Die Brunn-Minkowskische Ungleichung und ihr Spiegelbild sowie die isoperimetrische Eigenschaft der Kugel in der euklidischen und nichteuklidischen Geometrie. I*, Math. Nachr. **1** (1948), 81–157. MR 0028600 (10,471d)

MIT, Cambridge, MA 02139 USA
*E-mail address*: hcalpert@math.mit.edu
*E-mail address*: lguth@math.mit.edu